

EXPERT RESEARCH PROTOCOL

from 09/09/2015

Code: 15-08-20-1 (104q)
Customer:
Number of samples: Quantitative analysis of the sample
Methods: Agilent 1200, High-performance liquid chromatography (HPLC);
Column: Zorbax SB-C18 150 mm×2.1 mm, 3 mkm;
Detector – DAD, wavelength – 245 nm;
Detector – MSD, ionization method APCI Positive, SCAN (100 - 500 m/z)
Number of samples: 1
Subject: Clen (Clenbuterol)

1. The sample weighed 0.119 g is dissolved in 1 ml of solvent (MeOH). The extraction is 30 minutes in an ultrasonic bath.

2. 30 mg of the Standard (Clenbuterol, C2248000, European Pharmacopoeia Reference Standards, Ph. Eur.) dissolved in 1 ml of the solvent (Outlet 1-concentration-30 mg/ml).

3. Standard (Outlet 1) diluted 100 times (10 µl Outlet 1 + 990 µl of the solvent). Received Solution is Outlet 2 – concentration - 0.3 mg / ml (300 mcg / ml).

4. Standard (Outlet 2) diluted 6 times (200 µl Outlet 2+1000 µl of the solvent). Received Solution is Outlet 3 – concentration - 0.3 mg / ml (50 mcg / ml).

5. The Outlet 3 solution was used for making calibration dilutions (for injecting 1 µl of solution into the chromatograph):

T4 - Outlet 3 C = 50 mcg / ml;

T3 - 200 µl T4 + 200 µl of solvent C = 25 mcg / ml;

T2 - 200 µl T4 + 800 µl of solvent C = 10 mcg / ml;

T1 - 100 µl T4 + 900 µl of solvent C = 5 mcg / ml.

6. Research conditions:

Mobile phase: A - ACN-1% Formic acid (30%), B - H₂O-1% Formic acid (70%).

The elution mode is an isocratic.

The flow rate through the column: 0.3 ml/min. A thermostat: 30°C.

7. Single quadrupole mass analyzer is used for identification of the chemical elements. The samples were ionized in the chemical ionization mode at atmospheric pressure (APCI) with fixation of positive ions (Fig. 1)




Fig.1. The component output chromatogram of the sample, detector DA

8. The analysis results of the received peaks are shown on Fig. 2.




Fig.2. The analysis of the 1st peak

9. The received data of the analytical comparison results of MS and DA detectors with the calculated data on the test substance allow us to state that peak 1 refers to Clenbuterol.

10. The researched data of the standard samples for making a calibration chart is illustrated in the Table 1.

Tbl. 1. Peak values in accordance with the concentration of the test substance

Signal 1: DAD1 B, Sig=245,4 Ref=360,100

RetTime [min]	Lvl Sig	Amount [mkg/ml]	Area	Amt/Area	Ref Grp Name
2.007	1	5.00000	17.96171	2.78370e-1	Clenbuterol
		10.00000	34.09997	2.93255e-1	
		25.00000	89.43427	2.79535e-1	
		50.00000	170.09660	2.93951e-1	

11. The calibration chart (Fig. 3) is made form received data of the Table 1.




Fig.3. Calibration curve

12. The calculation of the concentration of the test sample from the HPLC data from the obtained calibration curve:

21.12 mcg/ml

21.32 mcg/ml

21.02 mcg/ml

21.48 mcg/ml

13. The calculated average is 21.24 mcg / ml.

14. The sample of the preparation of 0.119 g is dissolved in 1 ml of the solvent, therefore it will contain 21.24 mcg of the preparation.


15. Weight of the pills: 0.384 g, 0.363 g, 0.364 g, 0.383 g, 0.383 g. The average is 0.3754 g.

16. The amount of the sample in the pill is: 23.34 mcg.

Conclusion: Concentration of the test sample Clen (Clenbuterol) is 23.34 mcg in a pill.

Remarks:

1. This conclusion was issued based on the results of examination of the provided samples. This conclusion does not guarantee the overall quality of the products.
2. This conclusion can not be used for advertising purposes. It can not serve as an evidence in a lawsuit or to be used for other purposes.



AnabolicMenuStore